

Tiling the $(n^2, 1)$ -De Bruijn graph with n coassociative coalgebras ¹

Philippe LEROUX

Institut de Recherche Mathématique, Université de Rennes I and CNRS UMR 6625
Campus de Beaulieu, 35042 Rennes Cedex, France, philippe.leroux@univ-rennes1.fr

Abstract: We construct, via usual graph theory a class of associative dialgebras as well as a class of coassociative L -coalgebras, the two classes being related by a tool from graph theory called the line-extension. As a corollary, a tiling of the n^2 -De Bruijn graph with n (geometric supports of) coassociative coalgebras is obtained. We get, via the tiling of the $(3, 1)$ -De Bruijn graph, an example of cubical trialgebra defined by J-L Loday and M. Ronco. Other examples are obtained by letting $M_n(k)$ acts on the axioms defining such tilings. Examples of associative products which split into several associative ones are also given.

1 Introduction

The field k stands for the real field or the complex field. Moreover, all the vector spaces considered in this paper will have a finite or a denumerable basis. Let us recall the general setting of this article by summarising the main steps of our previous works [4, 5, 7, 6].

Definition 1.1 [Directed graph] A *directed graph* G is a quadruple [11], (G_0, G_1, s, t) where G_0 and G_1 are two denumerable sets respectively called the *vertex set* and the *arrow set*. The two mappings, $s, t : G_1 \rightarrow G_0$ are respectively called *source* and *terminus*. A vertex $v \in G_0$ is a *source* (resp. a *sink*) if $t^{-1}(\{v\})$ (resp. $s^{-1}(\{v\})$) is empty. A graph G is said *locally finite*, (resp. *row-finite*) if $t^{-1}(\{v\})$ is finite (resp. $s^{-1}(\{v\})$ is finite). Let us fix a vertex $v \in G_0$. Define the set $F_v := \{a \in G_1, s(a) = v\}$. A *weight* associated with the vertex v is a mapping $w_v : F_v \rightarrow k$. A directed graph equipped with a family of weights $w := (w_v)_{v \in G_0}$ is called a *weighted directed graph*.

In the sequel, directed graphs will be supposed locally finite and row finite. Let us introduce particular coalgebras named L -coalgebras ² and explain why this notion is interesting.

¹2000 *Mathematics Subject Classification*: 16W30; 05C20; 05C90. *Key words and phrases*: directed graphs, Hopf algebra, (Markov) L -coalgebra, coassociative co-dialgebra, cubical trialgebra, achirality.

²This notion has been introduced in [4] and developed in [4, 5, 7, 6].


Definition 1.2 [*L-coalgebra*] A *L-coalgebra* $(L, \Delta, \tilde{\Delta})$ over a field k is a k -vector space composed of a right part (L, Δ) , where $\Delta : L \rightarrow L^{\otimes 2}$, is called the right coproduct and a left part $(L, \tilde{\Delta})$, where $\tilde{\Delta} : L \rightarrow L^{\otimes 2}$, is called the left coproduct such that the coassociativity breaking equation, $(\tilde{\Delta} \otimes id)\Delta = (id \otimes \Delta)\tilde{\Delta}$, is verified. If $\Delta = \tilde{\Delta}$, the *L-coalgebra* is said *degenerate*. A *L-coalgebra* may have two counits, the right counit $\epsilon : L \rightarrow k$, verifying $(id \otimes \epsilon)\Delta = id$ and the left counit $\tilde{\epsilon} : L \rightarrow k$, verifying $(\tilde{\epsilon} \otimes id)\tilde{\Delta} = id$. A *L-coalgebra* is said *coassociative* if its two coproducts are coassociative. In this case the equation, $(\tilde{\Delta} \otimes id)\Delta = (id \otimes \Delta)\tilde{\Delta}$, is called the **entanglement equation** and we will say that its right part (L, Δ) is entangled to its left part $(L, \tilde{\Delta})$. Denote by τ , the *transposition* mapping, i.e. $L^{\otimes 2} \xrightarrow{\tau} L^{\otimes 2}$ such that $\tau(x \otimes y) = y \otimes x$, for all $x, y \in L$. The *L-coalgebra* L is said to be *L-cocommutative* if for all $v \in L$, $(\Delta - \tau\tilde{\Delta})v = 0$. A *L-bialgebra* (with counits $\epsilon, \tilde{\epsilon}$), is a *L-coalgebra* (with counits) and an unital algebra such that its coproducts and counits are homomorphisms. A *L-Hopf algebra*, H , is a *L-bialgebra* with counits equipped with right and left antipodes $S, \tilde{S} : H \rightarrow H$, such that: $m(id \otimes S)\Delta = \eta\epsilon$ and $m(\tilde{S} \otimes id)\tilde{\Delta} = \eta\tilde{\epsilon}$ or $m(S \otimes id)\Delta = \eta\epsilon$ and $m(id \otimes \tilde{S})\tilde{\Delta} = \eta\tilde{\epsilon}$.

Let $G = (G_0, G_1, s, t)$ be a directed graph equipped with a family of weights $(w_v)_{v \in G_0}$. Let us consider the free k -vector space kG_0 generated by G_0 . The set G_1 is then viewed as a subset of $(kG_0)^{\otimes 2}$ by identifying $a \in G_1$ with $s(a) \otimes t(a)$. The mappings source and terminus are then linear mappings still called source and terminus $s, t : (kG_0)^{\otimes 2} \rightarrow kG_0$, such that $s(u \otimes v) = u$ and $t(u \otimes v) = v$, for all $u, v \in G_0$. The family of weights $(w_v : F_v \rightarrow k)_{v \in G_0}$, is then viewed as a family of linear mappings. Let $v \in G_0$. Define the right coproduct $\Delta_M : kG_0 \rightarrow (kG_0)^{\otimes 2}$, such that $\Delta_M(v) := \sum_{i: a_i \in F_v} w_v(a_i) v \otimes t(a_i)$ and the left coproduct $\tilde{\Delta}_M : kG_0 \rightarrow (kG_0)^{\otimes 2}$, such that $\tilde{\Delta}_M(v) := \sum_{i: a_i \in P_v} w_{s(a_i)}(a_i) s(a_i) \otimes v$, where P_v is the set $\{a \in G_1, t(a) = v\}$. With these definitions the k -vector space kG_0 is a *L-coalgebra* called a finite Markov *L-coalgebra* since its coproducts Δ_M and $\tilde{\Delta}_M$ verify the coassociativity breaking equation $(\tilde{\Delta}_M \otimes id)\Delta_M = (id \otimes \Delta_M)\tilde{\Delta}_M$. This particular coalgebra is called in addition finite Markov (*L-coalgebra*) because for all $v \in G_0$, the sets F_v and P_v are finite and the coproducts have the form $\Delta_M(v) := v \otimes \dots$ and $\tilde{\Delta}_M(v) := \dots \otimes v$.

Assume we consider the Markov *L-coalgebra* just described and associate with each tensor product $\lambda u \otimes v$, where $\lambda \in k$ and $u, v \in G_0$, appearing in the definition of the coproducts, a directed arrow $u \xrightarrow{\lambda} v$. The weighted directed graph so obtained, called the *geometric support* of this *L-coalgebra*, is up to a graph isomorphism³, the directed graph we start with. Therefore, general *L-coalgebras* generalise the notion of weighted directed graph. If $(L, \Delta, \tilde{\Delta})$ is a *L-coalgebra* generated as a k -vector space by a set L_0 , then its geometric support $Gr(L)$ is a directed graph with vertex set $Gr(L)_0 = L_0$ and with arrow set $Gr(L)_1$, the set of tensor products $u \otimes v$, with $u, v \in L_0$, appearing in the definition of the coproducts of L . As a coassociative coalgebra is a particular *L-coalgebra*, we naturally construct its directed graph. We draw attention to the fact that a directed graph can be the geometric support of different *L-coalgebras*.

Example 1.3 The directed graph:

³A *graph isomorphism* $f : G \rightarrow H$ between two graphs G and H is a pair of bijection $f_0 : G_0 \rightarrow H_0$ and $f_1 : G_1 \rightarrow H_1$ such that $f_0(s_G(a)) = s_H(f_1(a))$ and $f_0(t_G(a)) = t_H(f_1(a))$ for all $a \in G_1$. All the directed graphs in this formalism will be considered up to a graph isomorphism.



is the geometric support associated with the degenerate L -coalgebra or coassociative coalgebra $\mathcal{F}$, generated by a, b, c and d , as a k -vector space, and described by the following coproduct: $\Delta a = a \otimes a + b \otimes c$, $\Delta b = a \otimes b + b \otimes d$, $\Delta c = d \otimes c + c \otimes a$, $\Delta d = d \otimes d + c \otimes b$ and the geometric support of the finite Markov L -coalgebra, generated by a, b, c and d , as a k -vector space, and described by the right coproduct: $\Delta_M a = a \otimes (a + b)$, $\Delta_M b = b \otimes (c + d)$, $\Delta_M c = c \otimes (a + b)$, $\Delta_M d = d \otimes (c + d)$ and the left coproduct: $\tilde{\Delta}_M a = (a + c) \otimes a$, $\tilde{\Delta}_M b = (a + c) \otimes b$, $\tilde{\Delta}_M c = (b + d) \otimes c$, $\tilde{\Delta}_M d = (b + d) \otimes d$.

Remark: Let $(\mathcal{G}, \Delta_M, \tilde{\Delta}_M)$ be a finite Markov L -coalgebra generated as a k -vector space by a set $\mathcal{G}_0$. If the family of weights $(w_v)_{v \in \mathcal{G}_0}$, $(\tilde{w}_v)_{v \in \mathcal{G}_0}$ used for describing right and left coproducts take values into $\mathbb{R}_+$ and if the right counit $\epsilon : v \mapsto 1$ exists, then the geometric support associated with $(\mathcal{G}, \Delta_M, \tilde{\Delta}_M)$ is a directed graph equipped with a family of probability vectors described by $(w_v)_{v \in \mathcal{G}_0}$.

Before going on, let us interpret what represents the L -cocommutativity in the case of a finite Markov L -coalgebra. A directed graph is said bi-directed if for any arrow from a vertex v_1 to a vertex v_2 , there exists an arrow from v_2 to v_1 . To take into account the bi-orientation of a directed graph in an algebraic way, we first embed this directed graph into the finite Markov L -coalgebra described above. We then notice that a directed graph is bi-directed if and only if $\Delta_M = \tau \tilde{\Delta}_M$. Therefore, in this algebraic framework, we are naturally led to consider the L -cocommutator $\ker(\Delta - \tau \tilde{\Delta})$. Dualizing this formula leads to consider an (L) -algebra D equipped with two products $\vdash$ and $\dashv$, verifying $(x \vdash y) \dashv z = x \vdash (y \dashv z)$, $x, y, z \in D$, and to consider the particular commutator $[x, y] := x \dashv y - y \vdash x$. The bracket, $[-, z]$, verifies the analogue of the “Jacobi identity”, called the Leibniz identity, i.e. $[[x, y], z] = [[x, z], y] + [x, [y, z]]$, if D is an algebra called an associative dialgebra [8].

Another motivation concerning associative dialgebras is the following. In a long-standing project whose ultimate aim is to study periodicity phenomena in algebraic K -theory, J-L. Loday in [8], and J-L. Loday and M. Ronco in [10] introduce several kind of algebras, one of which is the “non-commutative Lie algebras”, called *Leibniz algebras*. Such algebras D are described by a bracket $[-, z]$ verifying the so called Leibniz identity:

$$[[x, y], z] = [[x, z], y] + [x, [y, z]].$$

When the bracket is skew-symmetric, the Leibniz identity becomes the Jacobi identity and the Leibniz algebra turns out to be a Lie algebra. A way to construct such Leibniz algebra is to start with an *associative dialgebra*, that is a k -vector space D equipped with two associative products, $\vdash$ and $\dashv$, such that for all $x, y, z \in D$

1. $x \dashv (y \dashv z) = x \dashv (y \vdash z)$,
2. $(x \vdash y) \dashv z = x \vdash (y \dashv z)$,
3. $(x \dashv y) \vdash z = (x \vdash y) \vdash z$.

The associative dialgebra is then a Leibniz algebra by defining the bracket $[x, y] := x \dashv y - y \vdash x$, for all $x, y \in D$. The operad associated with associative dialgebras is then Koszul dual to the operad associated with dendriform algebras, a *dendriform algebra* Z being a k -vector space equipped with two binary operations, $\prec, \succ: Z \otimes Z \rightarrow Z$, satisfying the following axioms:

1. $(a \prec b) \prec c = a \prec (b \prec c) + a \prec (b \succ c)$,
2. $(a \succ b) \prec c = a \succ (b \prec c)$,
3. $(a \prec b) \succ c + (a \succ b) \succ c = a \succ (b \succ c)$.

This notion dichotomizes the notion of associativity since the product $a * b = a \prec b + a \succ b$, for all $a, b \in Z$ is associative. Otherwise stated, the associative product $*$ splits into two operations $\prec$ and $\succ$.

The first important result of this paper is the construction, via Markov L -coalgebras, i.e. via usual graph theory of a class of L -cocommutative and coassociative codialgebras as well as a class of coassociative L -coalgebras, the two classes being related by a tool from graph theory called the line-extension. The second one is the construction of a tiling of the $(n^2, 1)$ -De Bruijn graph with n (geometric supports of) coassociative coalgebras. As a corollary, we obtain examples of cubical trialgebras, a notion developed by J-L Loday and M. Ronco in [10] and splittings of associative products into several associative ones.

Let us briefly introduce the organisation of the paper. In section 2, we display the notion of coassociative co-dialgebras and recall the definition of the De-Bruijn graphs. We prove that the $(2, 1)$ -De Bruijn graph, viewed as a Markov co-dialgebra, yields, by line-extension, the geometric support of the coassociative coalgebra $\mathcal{F}$, generated by say a, b, c and d , as a k -vector space, and described by the following coproduct: $\Delta a = a \otimes a + b \otimes c$, $\Delta b = a \otimes b + b \otimes d$, $\Delta c = d \otimes c + c \otimes a$, $\Delta d = d \otimes d + c \otimes b$. Inspired by a previous work [7], relationships between the Markovian coproduct of the $(2, 1)$ -De Bruijn graph and the coassociative coproduct of $\mathcal{F}$ are also given. In section 3, we prove the existence of another structure associated with $\mathcal{F}$. This structure is also a coassociative coalgebra, called by convention the left structure of $\mathcal{F}$. As the two structures of $\mathcal{F}$ are entangled by the entanglement equation, we conclude that $\mathcal{F}$ is a coassociative L -coalgebra. Moreover, gluing their two associated geometric supports yields the $(4, 1)$ -De Bruijn graph. Since the intersection of the two arrow sets of their geometric supports is empty, we assert that the $(4, 1)$ -De Bruijn graph can be tiled by the two geometric supports of the coassociative L -coalgebra $\mathcal{F}$.

The consequences of such a left structure on $Sl_q(2)$ are also explored. If $\mathcal{F}$ stands for $Sl_q(2)$, we notice that the usual algebraic relations of the Hopf algebra $Sl_q(2)$ do not embed this new structure into a bialgebra. However, it allows us to construct a map which verify the same axioms as an antipode map. We yield also a left structure for $SU_q(2)$.

This work ends, in section 4, by showing that the $(n^2, 1)$ -De Bruijn graph can also be tiled with n (geometric supports of) coassociative coalgebras. As a consequence, we give examples of cubical trialgebras and put forward an important notion which is the achirality of a L -coalgebra. we give also Leibniz co-derivative on these directed graphs.

2 On the De Bruijn graph families

The description of weighted directed graphs and coassociative coalgebras can be embedded into the L -coalgebra framework. In [4], we establish an important theorem.

Proposition 2.1 *Any coassociative coalgebra (C, Δ_C) , with a group-like element can be embedded into a non degenerate L -coalgebra.*

Proof: Let (C, Δ_C) be a coassociative coalgebra. Suppose e is a group-like element, i.e. $\Delta_C e = e \otimes e$. Define the coassociative coproducts $\delta(c) := c \otimes e$ and $\tilde{\delta}(c) := e \otimes c$ for all $c \in C$. Define also the linear map $\vec{d} : C \rightarrow C \otimes C$ such that $\vec{d}(c) = \Delta_C - \delta_f(c)$ and the linear map $\overleftarrow{d} : C \rightarrow C \otimes C$ such that $\overleftarrow{d}(c) = \Delta_C - \tilde{\delta}_f(c)$. These linear maps, $\overleftarrow{d}$ and $\vec{d}$, turn the coassociative coalgebra (C, Δ) into a non degenerate L -coalgebra $(C, \vec{d}, \overleftarrow{d})$. $\square$

Let C be a bialgebra with unit e . The two new coproducts, $\overleftarrow{d}, \vec{d} : C \rightarrow C^{\otimes 2}$, turn out to be Ito derivative⁴. Starting with a Markov L -coalgebra $\mathcal{G}$, we can recover a similar theorem. However, whereas in the coassociative coalgebra case, these new two coproducts map “the vertex set” C into the “arrow sets” $C^{\otimes 2}$, the new coproducts, $\overleftarrow{d}_{\mathcal{G}}$ and $\vec{d}_{\mathcal{G}}$, of a Markov L -coalgebra $\mathcal{G}$ map “the arrow set” $C^{\otimes 2}$ into the “paths of length 2” modelised by $C^{\otimes 3}$. This observation suggests the necessity of studying the line-extension of directed graphs, a notion defined in the sequel.

2.1 Line-extension of directed graphs

The aim of this subsection is to show that some De Bruijn directed graphs, seen as Markov L -coalgebras are also coassociative co-dialgebras and that the line-extension of these directed graphs can be viewed as geometric supports associated with some well-known coassociative coalgebras. We start with two definitions.

Definition 2.2 [De Bruijn graph] A (p, n) De-Bruijn sequence on the alphabet $\Sigma = \{a_1, \dots, a_p\}$ is a sequence $(s_1, \dots, s_m)$ of $m = p^n$ elements $s_i \in \Sigma$ such that subsequences of length n of the form $(s_i, \dots, s_{i+n-1})$ are distinct, the addition of subscripts being done modulo m . A (p, n) -De Bruijn graph is a directed graph whose vertices correspond to all possible strings $s_1 s_2 \dots s_n$ of n symbols from Σ . There are p arcs leaving the vertex $s_1 s_2 \dots s_n$ and leading to the adjacent node $s_2 s_3 \dots s_n \alpha$, $\alpha \in \Sigma$. Therefore the $(p, 1)$ -De Bruijn graph is the directed graph with p vertices, complete, with a loop at each vertex.

Fix $n \geq 1$. Let $D_{(n,1)} = ((D_{(n,1)})_0, (D_{(n,1)})_1, s, t)$ be the $(n, 1)$ -De Bruijn graph. Its natural Markov L -coalgebra is defined as follows. Denote by v_i , $1 \leq i \leq n$, the vertices of $D_{(n,1)}$. Embed $(D_{(n,1)})_0$ into its free k -vector space. Define the coproducts $\Delta_M, \tilde{\Delta}_M : k(D_{(n,1)})_0 \rightarrow k(D_{(n,1)})_0^{\otimes 2}$, such that for all i , $\Delta_M v_i := v_i \otimes \sum_j v_j$ and $\tilde{\Delta}_M v_i := \sum_j v_j \otimes v_i$. There are obvious left and right counits, $\tilde{\epsilon}(v_i) = \epsilon(v_i) = \frac{1}{n}$, for all $v_i \in (D_{(n,1)})_0$.

⁴Let A be an associative algebra with unit e and $f : A \rightarrow A$ be a linear map. The map f is said to be an Ito derivative if $f(e) = 0$ and $f(xy) = f(x)f(y) + xf(y) + f(x)y$, for all $x, y \in A$.

Definition 2.3 [Line-extension] The *line-extension* of a directed graph G , with vertex set $G_0 = \{j_1, \dots, j_n\}$ and arrow set $G_1 \subseteq G_0 \times G_0$, denoted by $E(G)$, is the directed graph with vertex set $E(G)_0 := G_1$ and the arrow set $E(G)_1 \subseteq G_1 \times G_1$ defined by $((j_k, j_l), (j_e, j_f))$ iff $j_l = j_e$.

The notion of associative dialgebra, due to J-L. Loday [8], is a notion which generalizes the notion of algebra. Associative dialgebras, via the notion of dendriform algebra ⁵, are closely related to (planar binary) trees, which became an important tool in quantum field theory [3, 2]. Here, we are interested in the notion of coassociative co-dialgebra.

Definition 2.4 [Coassociative co-dialgebra of degree n] Motivated by line-extension of (geometric support of) Markov L -coalgebra, we define in [4], (Markov) L -coalgebra of degree n , $n > 0$. Similarly, let Δ_n and $\tilde{\Delta}_n$ be two n -linear mappings $D^{\otimes n} \rightarrow D^{\otimes n+1}$, where D is a k -vector space. The k -vector space $(D, \Delta_n, \tilde{\Delta}_n)$ is said to be a *coassociative co-dialgebra* of degree n if the following axioms are verified:


1. Δ_n and $\tilde{\Delta}_n$ are coassociative,
2. $(id \otimes \Delta_n)\Delta_n = (id \otimes \tilde{\Delta}_n)\Delta_n$,
3. $(\tilde{\Delta}_n \otimes id)\tilde{\Delta}_n = (\Delta_n \otimes id)\tilde{\Delta}_n$,
4. $(\tilde{\Delta}_n \otimes id)\Delta_n = (id \otimes \Delta_n)\tilde{\Delta}_n$.

As in the L -coalgebra case, such a space may have a right counit $\epsilon_n : D^{\otimes n} \rightarrow D^{\otimes n-1}$ such that: $(id \otimes \epsilon_n)\Delta_n = id$ and a left counit $\tilde{\epsilon}_n : D^{\otimes n} \rightarrow D^{\otimes n-1}$ such that: $(\tilde{\epsilon}_n \otimes id)\tilde{\Delta}_n = id$. By convention $D^{\otimes 0} := k$. A coassociative co-dialgebra of degree 1 will be also called a coassociative co-dialgebra.

Proposition 2.5 *A k -vector space C equipped with two coproducts δ_f and $\tilde{\delta}_f$ such that $\delta_f(c) = c \otimes e$ and $\tilde{\delta}_f(c) = e \otimes c$, for all $c \in C$ is a Markov coassociative codialgebra.*

Proof: Straightforward since $\delta_f(e) = \tilde{\delta}_f(e)$. □

Example 2.6 [The flower graph] A unital algebra, A with unit 1, carries a non trivial Markov L -bialgebra, obtained from the equality, $(1 \cdot a) \cdot 1 = 1 \cdot (a \cdot 1)$. The coproducts are $\delta_f(a) = a \otimes 1$ and $\tilde{\delta}_f(a) = 1 \otimes a$, $a \in A$. Its geometric support is called the flower graph.



Example of geometric support associated with an algebra $k\langle a, b, c, d \rangle \oplus k1$.

⁵The operads Dias (associative dialgebra) and Dend (dendriform algebra) are dual in the operad sense, see [8].

Theorem 2.7 Any coassociative coalgebra (C, Δ) , (respectively bialgebra, Hopf algebra), can be embedded into a L -coalgebra of degree n , (respectively, L -bialgebra of degree n , L -Hopf algebra ⁶ of degree n), with $n > 1$.

Proof: Fix $n > 1$. Consider Δ , the coproduct of such a coassociative coalgebra C . Set: $id_n := \underbrace{id \otimes id \otimes \dots \otimes id}_n$ and $\Delta_n := id_{n-1} \otimes \Delta$ and $\tilde{\Delta}_n := \Delta \otimes id_{n-1}$. The right and left coassociative coproducts Δ_n and $\tilde{\Delta}_n$ map $C^{\otimes n}$ into $C^{\otimes n+1}$ and the entanglement equation is realised. The right and left counits are $\epsilon_n := id_{n-1} \otimes \epsilon$ and $\tilde{\epsilon}_n := \epsilon \otimes id_{n-1}$, since for instance $(id \otimes \epsilon_n)\Delta_n = id_n$. If the counit and the coproduct of C are unital homomorphisms, so are the new coproducts and counits. If C is a Hopf algebra with antipode s , we embed C into a L -Hopf algebra of degree n since $(id_{n-1} \otimes m)(id_n \otimes s)\Delta_n = \eta_n \epsilon_n$ and $(m \otimes id_{n-1})(s \otimes id_n)\tilde{\Delta}_n = \tilde{\eta}_n \tilde{\epsilon}_n$. $\square$

Theorem 2.8 If $n = 2$, any coassociative coalgebra (C, Δ) , can be viewed as a coassociative co-dialgebra of degree 2.

Proof: Straightforward by using the definition of Δ_2 and the fact that Δ is coassociative. $\square$

Proposition 2.9 Let $D_{(n,1)} = ((D_{(n,1)})_0, (D_{(n,1)})_1)$, be the $(n, 1)$ -De Bruijn graph and consider its free k -vector space $k(D_{(n,1)})_0$. The Markovian coproducts of the Markov L -coalgebra $k(D_{(n,1)})_0$ associated with the $(n, 1)$ -De Bruijn graph define a coassociative co-dialgebra.

Proof: Straightforward. $\square$

Proposition 2.10 There exists a coassociative coalgebra whose geometric support is the line-extension of the $(n, 1)$ -De Bruijn graph.

Proof: Let us denote the arrow emerging from a given vertex v_i to a vertex v_j , with $i, j = 1, \dots, n$ of the $(n, 1)$ -De Bruijn graph $D_{(n,1)}$ by a_{ij} . The new vertices of $E(D_{(n,1)})$ are denoted by a_{ij} and the arrows are denoted by $((il), (lj))$. Consider the free k -vector space generated by the set $E(D_{(n,1)})_0 := \{a_{ij}; i, j = 1 \dots n\}$ and define $\Delta a_{ij} = \sum_l a_{il} \otimes a_{lj}$, this coproduct is coassociative and the geometric support associating with the coassociative coalgebra $(kE(D_{(n,1)})_0, \Delta)$ is easily seen to be $E(D_{(n,1)})$. It has an obvious counit, $a_{ij} \mapsto 0$ if $i \neq j$ and $a_{ij} \mapsto 1$ otherwise. $\square$

Corollary 2.11 Recall that $\mathcal{F}$ is generated by a, b, c and d , as a k -vector space, and described by the following coproduct: $\Delta a = a \otimes a + b \otimes c$, $\Delta b = a \otimes b + b \otimes d$, $\Delta c = d \otimes c + c \otimes a$, $\Delta d = d \otimes d + c \otimes b$. The line-extension of the $(2, 1)$ -De Bruijn graph can be equipped with the coassociative coproduct associated with the coalgebra $\mathcal{F}$.

⁶Set $id_n := \underbrace{id \otimes id \otimes \dots \otimes id}_n$, $n > 0$. A L -Hopf algebra of degree n , $(H, \Delta_H, \tilde{\Delta}_H)$, is by definition a L -bialgebra

of degree n , equipped with right and left counits, $\tilde{\epsilon}_H, \epsilon_H$ of degree n , such that its antipodes $S, \tilde{S} : H \rightarrow H$ verify $(id_{n-1} \otimes m)(id_n \otimes S)\Delta_H = \eta_n \epsilon_H$ and $(m \otimes id)(\tilde{S} \otimes id_n)\tilde{\Delta}_H = \tilde{\eta}_n \tilde{\epsilon}_H$, with $\eta_n, \tilde{\eta}_n : H^{\otimes(n-1)} \rightarrow H^{\otimes n}$ such that $\eta_n(h) := h \otimes 1_H$ and $\tilde{\eta}_n(h) := 1_H \otimes h$, $h \in H^{\otimes(n-1)}$.

Proposition 2.12 *Let B be a (Markov) coassociative co-dialgebra with coproducts $\tilde{\Delta}$ and Δ and C be a coassociative coalgebra with coproduct Δ_C . Then $B \otimes C$ is a coassociative co-dialgebra with coproduct $\delta_{B \otimes C} := (id \otimes \tau \otimes id)\Delta \otimes \Delta_C$ and $\tilde{\delta}_{B \otimes C} := (id \otimes \tau \otimes id)\tilde{\Delta} \otimes \Delta_C$. Similarly, $C \otimes B$ is a coassociative co-dialgebra with $\delta_{C \otimes B} := (id \otimes \tau \otimes id)\Delta_C \otimes \Delta$ and $\tilde{\delta}_{C \otimes B} := (id \otimes \tau \otimes id)\Delta_C \otimes \tilde{\Delta}$.*

Proof: Straightforward. $\square$

We end this section on De-Bruijn graphs and coassociative codialgebras, by constructing another family of coassociative co-dialgebra. The idea is to give the $(n, 1)$ -De Bruijn graph an attractor rôle.

Proposition 2.13 *Let n and m be two integers different from zero. Let D be the k -vector space $k\langle x_1, \dots, x_m \rangle \oplus k\langle \alpha_1, \dots, \alpha_n \rangle$. Define for all $i = 1, \dots, m$, $\Delta x_i = \sum_{j=1}^n x_i \otimes \alpha_j$, $\tilde{\Delta} x_i = \sum_{j=1}^n \alpha_j \otimes x_i$ and for all $j = 1, \dots, n$, $\Delta \alpha_j = \sum_{p=1}^n \alpha_j \otimes \alpha_p$, $\tilde{\Delta} \alpha_j = \sum_{p=1}^n \alpha_p \otimes \alpha_j$. These Markovian coproducts embed D into a coassociative co-dialgebra.*

Proof: Straightforward. $\square$

Remark: Consider the k -vector space of linear maps $L(D, A)$ which map $(D, \Delta, \tilde{\Delta})$, a coassociative coalgebra into an associative algebra A equipped with a product m . The space $L(D, A)$ is then embedded into an associative dialgebra by defining for all $f, g \in L(D, A)$, the two convolution products: $f \dashv g := m(f \otimes g)\Delta$ and $f \vdash g := m(f \otimes g)\tilde{\Delta}$.

2.2 Relationships between the $(2, 1)$ -De Bruijn L -coalgebra and $\mathcal{F}$

Motivated by a previous work [7], the aim of this subsection is to study the relationships between the $(2, 1)$ -De Bruijn graph, seen as a Markov L -coalgebra and its line-extension seen as the (geometric support of the) coassociative coalgebra $\mathcal{F}$. Let X, Y be two non commuting operators, we consider the non commutative algebra $A = k\langle X, Y \rangle \oplus k1$, with $XY = \eta YX$ and $\eta \in k \setminus \{0\}$. We equip A with the following Markovian coproducts $\Delta_M X = X \otimes X + X \otimes Y$ and $\Delta_M Y = Y \otimes X + Y \otimes Y$, $\tilde{\Delta}_M X = Y \otimes X + X \otimes X$, $\tilde{\Delta}_M Y = X \otimes Y + Y \otimes Y$ and set $\tilde{\Delta}_M 1 = \Delta_M 1 = 1 \otimes 1$. Set $a = X \otimes X$, $b = X \otimes Y$, $c = Y \otimes X$, $d = Y \otimes Y$. We would like to find operators which give the coassociative coproduct of $\mathcal{F}$ from the coassociative co-dialgebra A . For that, we define:

$$\overrightarrow{\Delta}_M : A \rightarrow A^{\otimes 2} \times A^{\otimes 2}, \quad X \mapsto \overrightarrow{\Delta}_M(X) = \begin{pmatrix} X \otimes X \\ X \otimes Y \end{pmatrix}, \quad Y \mapsto \overrightarrow{\Delta}_M(Y) = \begin{pmatrix} Y \otimes Y \\ Y \otimes X \end{pmatrix}, \quad 1 \mapsto \overrightarrow{\Delta}_M(1) = \begin{pmatrix} 1 \otimes 1 \\ 1 \otimes 1 \end{pmatrix},$$

and

$$\overleftarrow{\Delta}_M : A \rightarrow A^{\otimes 2} \times A^{\otimes 2}, \quad X \mapsto \overleftarrow{\Delta}_M(X) = \begin{pmatrix} X \otimes X \\ Y \otimes X \end{pmatrix}, \quad Y \mapsto \overleftarrow{\Delta}_M(Y) = \begin{pmatrix} Y \otimes Y \\ X \otimes Y \end{pmatrix}, \quad 1 \mapsto \overleftarrow{\Delta}_M(1) = \begin{pmatrix} 1 \otimes 1 \\ 1 \otimes 1 \end{pmatrix}.$$

If we define the bilinear map $\langle \cdot ; \cdot \rangle_* : (A^{\otimes 2} \times A^{\otimes 2})^{\times 2} \rightarrow A^{\otimes 2}$, $((z_1, z_2), (z_3, z_4)) \mapsto z_1 z_3 + z_2 z_4$, we recover $\Delta_M(X) = \langle \overrightarrow{\Delta}_M(X), \overrightarrow{\Delta}_M(1) \rangle_*$ and $\Delta_M(Y) = \langle \overrightarrow{\Delta}_M(Y), \overrightarrow{\Delta}_M(1) \rangle_*$. Define the bilinear map $\square : A^{\otimes 2} \times A^{\otimes 2} \rightarrow A^{\otimes 2} \otimes A^{\otimes 2}$ such that:

$$(y_1 \otimes y_2) \square (y_3 \otimes y_4) := (id \otimes \tau \otimes id)((y_1 \otimes y_2) \otimes (y_3 \otimes y_4)) := (y_1 \otimes y_3) \otimes (y_2 \otimes y_4).$$

Define also $\langle \cdot ; \cdot \rangle$ and $\text{per} : (A^{\otimes 2} \times A^{\otimes 2})^{\times 2} \rightarrow A^{\otimes 2} \otimes A^{\otimes 2}$ such that $((z_1, z_2), (z_3, z_4)) \mapsto z_1 \square z_3 + z_2 \square z_4$ and the “permanent” $\text{per}((z_1, z_2), (z_3, z_4)) := z_1 \square z_4 + z_2 \square z_3$. Let us express the relations between Markovian coproducts of the $(2, 1)$ -De Bruijn graph and the coassociative coproduct of $\mathcal{F}$.

Proposition 2.14 *The relations are: $\langle \vec{\Delta}_M(X), \vec{\Delta}_M(X) \rangle = \Delta(a)$, $\langle \vec{\Delta}_M(Y), \vec{\Delta}_M(Y) \rangle = \Delta(d)$, $\text{per}(\vec{\Delta}_M(X), \vec{\Delta}_M(Y)) = \Delta(b)$, $\text{per}(\vec{\Delta}_M(Y), \vec{\Delta}_M(X)) = \Delta(c)$.*

Proof: Straightforward. For instance, $\langle \vec{\Delta}_M(X), \vec{\Delta}_M(X) \rangle = (X \otimes X) \square (X \otimes X) + (X \otimes Y) \square (Y \otimes X) = a \otimes a + b \otimes c = \Delta(a)$. $\square$


We can also recover algebraic relations of $Sl_q(2)$, except the q -determinant which is equal to zero, i.e. $ad - q^{-1}bc = 0$ instead of one. For checking the algebraic relations, we contract the arrows a, b, c, d into the vertices $\bar{a} = X^2, \bar{b} = XY, \bar{c} = YX, \bar{d} = Y^2$, thanks to the usual product of A , (recall that such a product maps the arrow set $A^{\otimes 2}$ into the vertex set A).

Proposition 2.15 *With $XY = \eta YX$, we obtain $\bar{a}\bar{b} = \eta^2 \bar{b}\bar{a}$, $\bar{c}\bar{b} = \bar{b}\bar{c}$, $\bar{a}\bar{c} = \eta^2 \bar{c}\bar{a}$, $\bar{a}\bar{d} = \eta^2 \bar{b}\bar{c}$, $\bar{b}\bar{d} = \eta^2 \bar{d}\bar{b}$, $\bar{c}\bar{d} = \eta^2 \bar{d}\bar{c}$, $\bar{a}\bar{d} - \bar{d}\bar{a} = (\eta^2 - \eta^{-2})\bar{b}\bar{c}$, $\bar{a}\bar{d} - \eta^2 \bar{b}\bar{c} = 0$.*

Proof: For instance, $\bar{a}\bar{b} = XXXY = \eta^2 XYXX = \eta^2 \bar{b}\bar{a}$, $\bar{c}\bar{d} = YXY Y = \eta^2 YYYX = \eta^2 \bar{d}\bar{c}$, $\bar{a}\bar{d} = XXY Y = \eta^2 XY YX = \eta^2 \bar{b}\bar{c}$, and so on. By setting $\eta^2 = q^{-1}$, we recover the usual algebraic relations for $Sl_q(2)$, except the q -determinant which is equal to 0. $\square$

3 The left part of $\mathcal{F}$

The aim of this section is to prove that there exists a coassociative coalgebra $(\mathcal{F}, \tilde{\Delta})$ entangled to the usual coassociative coalgebra $(\mathcal{F}, \Delta)$, i.e. to prove that $(\mathcal{F}, \Delta, \tilde{\Delta})$ is a coassociative L -coalgebra. By convention, we call $(\mathcal{F}, \tilde{\Delta})$, the left part of $(\mathcal{F}, \Delta, \tilde{\Delta})$. Its associated geometric support is:



Geometric support associated with the left part $(\mathcal{F}, \tilde{\Delta})$.

This structure can be obtained, for instance, by inverting the map $\langle \cdot, \cdot \rangle$ and the map per in the equations defining the usual coproduct Δ of $\mathcal{F}$ obtained from the $(2, 1)$ -De Bruijn graph in proposition 2.14. The relations for the new coproduct $\tilde{\Delta}$ are,

$$\tilde{\Delta}b = b \otimes b + a \otimes d, \quad \tilde{\Delta}c = c \otimes c + d \otimes a, \quad \tilde{\Delta}a = b \otimes a + a \otimes c, \quad \tilde{\Delta}d = c \otimes d + d \otimes b.$$

Remark: In [4], we defined a matrix product $U\bar{\otimes}W$, where U, W are two matrices and $(U\bar{\otimes}W)_{ij} := \sum_k U_{ik} \otimes W_{kj}$. With:

$$U = \begin{pmatrix} a & b \\ c & d \end{pmatrix}, \quad \tilde{U} = \begin{pmatrix} b & a \\ d & c \end{pmatrix},$$

the operation, $\sim$, meaning here the permutation of the two columns, the definition of the two coproducts can be recovered. Indeed, $(U\bar{\otimes}U)_{ij} := \sum_k U_{ik} \otimes U_{kj} := \Delta U_{ij}$ yields the right coproduct and the left coproduct $\tilde{\Delta}$ can be recovered by computing $\tilde{U}\bar{\otimes}\tilde{U}$.

Remark: The linear map $\tilde{\epsilon} : \mathcal{F} \rightarrow k$, such that:

$$\tilde{\epsilon}(a) = \tilde{\epsilon}(d) = 0, \quad \tilde{\epsilon}(b) = \tilde{\epsilon}(c) = 1,$$

is a counit map for $\tilde{\Delta}$, i.e. $(\tilde{\epsilon} \otimes id)\tilde{\Delta} = (id \otimes \tilde{\epsilon})\tilde{\Delta} = id$.

Theorem 3.1 *The new coproduct $\tilde{\Delta}$ is coassociative and verifies the entanglement equation $(\tilde{\Delta} \otimes id)\Delta = (id \otimes \Delta)\tilde{\Delta}$. Moreover the two coproducts verify also $(\Delta \otimes id)\tilde{\Delta} = (id \otimes \tilde{\Delta})\Delta$.*

Proof: The coassociativity is straightforward. The proof of the two equations is also straightforward by using the matrix product defined above. $\square$


Remark: The axioms of coassociative co-dialgebras are not satisfied.

Definition 3.2 [Chiral, Achiral] Let $(L, \Delta, \tilde{\Delta})$ be a coassociative L -coalgebra with right coproduct Δ and left coproduct $\tilde{\Delta}$. The k -vector space $(L, \Delta, \tilde{\Delta})$ is *chiral* iff $(\tilde{\Delta} \otimes id)\Delta = (id \otimes \Delta)\tilde{\Delta}$ and $(\Delta \otimes id)\tilde{\Delta} \neq (id \otimes \tilde{\Delta})\Delta$. This means that the entanglement equation “differentiates” the left part $(L, \tilde{\Delta})$ from the right part (L, Δ) . To the contrary, $(L, \Delta, \tilde{\Delta})$ is said to be *achiral* iff $(\tilde{\Delta} \otimes id)\Delta = (id \otimes \Delta)\tilde{\Delta}$ and $(\Delta \otimes id)\tilde{\Delta} = (id \otimes \tilde{\Delta})\Delta$, i.e. the axioms of an achiral coassociative L -coalgebra are globally invariant under the permutation $\tilde{\Delta} \leftrightarrow \Delta$. More generally, an algebra $(A, \bullet_1, \dots, \bullet_n)$ equipped with n operations $\bullet_1, \dots, \bullet_n : A^{\otimes 2} \rightarrow A$, verifying axioms $AX_1, \dots, AX_p$ is said to be achiral if $(A, \bullet_{\sigma(1)}, \dots, \bullet_{\sigma(n)})$, where σ is a permutation, verifies also the same axioms, i.e. the axioms $AX_1, \dots, AX_p$ are globally invariant under the action of any permutations σ . The dualisation of this definition is straightforward.

Remark: In the theorem 3.1, the entanglement equation is verified, even by inverting the rôle of the left and right coproducts. We say that $(\mathcal{F}, \Delta, \tilde{\Delta})$ is an achiral L -coalgebra, since the left and right parts are entangled by the entanglement equation which do not differentiate them. To the contrary, for instance, observe that the two coproducts δ_f and $\tilde{\delta}_f$ associated with an unital associative algebra, embed the algebra into a chiral L -bialgebra, see also [5].

Theorem 3.3 *The two geometric supports associated with the left part $(\mathcal{F}, \tilde{\Delta})$ and the right part $(\mathcal{F}, \Delta)$ of the coassociative L -coalgebra $(\mathcal{F}, \Delta, \tilde{\Delta})$, obtained from the line-extension of the $(2, 1)$ -De Bruijn graph, glued together yield the $(4, 1)$ -De Bruijn graph. Moreover the intersection of their arrow sets is empty.*

Proof: [The gluing of the left and right parts of $(\mathcal{F}, \Delta, \tilde{\Delta})$] When we glue the geometric support associated with $(\mathcal{F}, \Delta)$ with its left part $Gr(\mathcal{F}, \tilde{\Delta})$, we obtain the $(4, 1)$ -De Bruijn directed graph.



The (4, 1)-De Bruijn graph.

The intersection of the arrow sets of these two graphs is empty. $\square$

The (4, 1)-De Bruijn graph is tiled with two (geometric supports of) entangled coassociative coalgebras, the entanglement being achiral.

3.1 Consequences

3.1.1 An example of achiral L -Hopf algebra

Proposition 3.4 *If $(\mathcal{F}, \Delta, \tilde{\Delta})$ is also a unital algebra and a, b, c, d commute pairwise. Then $(\mathcal{F}, \Delta, \tilde{\Delta})$ will be an achiral L -bialgebra. Moreover, if $ad - bc = 1$, define the linear map $S_{\mathcal{F}}$ which maps $a \mapsto d$, $d \mapsto a$, $b \mapsto -b$ and $c \mapsto -c$ and the linear map $\tilde{S}_{\mathcal{F}}$ which maps $b \mapsto -c$, $c \mapsto -b$, $a \mapsto a$, $d \mapsto d$, then the right (resp. left) part of $(\mathcal{F}, \Delta, \tilde{\Delta})$ is a Hopf algebra, i.e. $(\mathcal{F}, \Delta, \tilde{\Delta})$ is an achiral L -Hopf algebra.*

Proof: Straightforward. $\square$

3.1.2 An example of chiral L -bialgebra

Consider the algebra generated by x, p and g such that $pg = gp$, $xg = \eta gx$, $xp = \mu px$, with η, μ two scalars different from zero. Define the (left) part by: $\tilde{\Delta}x = x \otimes p + g \otimes x$, $\tilde{\Delta}p = p \otimes p$, $\tilde{\Delta}g = g \otimes g$ with counit $\tilde{\epsilon}: x \mapsto 0$ and $p, g \mapsto 1$. Define the (right) part by: $\Delta x = x \otimes x$, $\Delta p = p \otimes x$, $\Delta g = g \otimes x$.

Proposition 3.5 *The two coproducts are coassociative and verify the entanglement equation $(id \otimes \Delta)\tilde{\Delta} = (\tilde{\Delta} \otimes id)\Delta$. Moreover the two coproducts Δ , $\tilde{\Delta}$ and the counit $\tilde{\epsilon}$ are homomorphisms.*

Proof: Tedious but straightforward by noticing that it is a consequence of the previous subsection on the left part of $\mathcal{F}$, by setting formally $b = 0$. For instance, we check the entanglement equation on x . We get, $x \xrightarrow{\Delta} x \otimes x \xrightarrow{\tilde{\Delta} \otimes id} x \otimes p \otimes x + g \otimes x \otimes x$ and $x \xrightarrow{\tilde{\Delta}} x \otimes p + g \otimes x \xrightarrow{id \otimes \Delta} x \otimes p \otimes x + g \otimes x \otimes x$. For the homomorphism property, we get for instance $\Delta xg = \Delta x \cdot \Delta g = xg \otimes xx$ and $\Delta gx = \Delta g \cdot \Delta x = gx \otimes xx$, thus $\Delta xg = \eta \Delta gx$ and so forth. $\square$

3.1.3 Consequences for $Sl_q(2)$

Proposition 3.6 *With the usual algebraic relations of the Hopf algebra $Sl_q(2)$, whose usual product will be denoted by m , the linear map $\tilde{S} : Sl_q(2) \rightarrow Sl_q(2)$ defined by $b \mapsto -q^{-1}c$; $c \mapsto -qb$; $a \mapsto a$; $d \mapsto d$, verifies $m(id \otimes \tilde{S})\tilde{\Delta} = m(\tilde{S} \otimes id)\tilde{\Delta} = 1 \cdot \tilde{\epsilon}$. Moreover, the map $\tilde{S}$ can be extended to an unital algebra map. As $1, a, d$ and bc are fixed points of $\tilde{S}$, the q -determinant of the matrix U is invariant by applying $\tilde{S}$.*

Proof: Let us check the antipode property. For that, multiply $\tilde{U}$ by the following matrix

$$\tilde{S}(\tilde{U}) = \begin{pmatrix} -q^{-1}c & a \\ d & -qb \end{pmatrix},$$

which is the matrix obtained from $\tilde{U}$ by the left antipode $\tilde{S}$. We compute

$$\widetilde{\tilde{U}\tilde{S}(\tilde{U})} = \begin{pmatrix} ba - qab & ad - q^{-1}bc \\ da - qcb & cd - q^{-1}dc \end{pmatrix}.$$

This matrix must be equal to:

$$\tilde{\epsilon}(U) = \begin{pmatrix} \tilde{\epsilon}(a) = 0 & \tilde{\epsilon}(b) = 1 \\ \tilde{\epsilon}(c) = 1 & \tilde{\epsilon}(d) = 0 \end{pmatrix},$$

which is the case since we know that $da = 1 + qbc$. We obtain the same result if we consider $\widetilde{\tilde{S}(\tilde{U})\tilde{U}}$. The extension of the definition of $\tilde{S}$ to a homomorphism is straightforward. $\square$

Remark: The algebraic relations of $Sl_q(2)$ do not embed its left part into a bialgebra. This fact will be a motivation for introducing coassociative manifolds in [5].

3.1.4 The left part of $SU_q(2)$

A consequence of what was done for the coassociative coalgebra $\mathcal{F}$ is the existence of a left part for the Hopf algebra $SU_q(2)$. We recall that $SU_q(2)$ is an $*$ -algebra with two generators, a and c such that $ac = qca$, $ac^* = qc^*a$, $cc^* = c^*c$, $a^*a + c^*c = 1$, $aa^* + q^2cc^* = 1$, with q a real different from 0. The coproduct is given by $\Delta_1 a = a \otimes a - qc^* \otimes c$ and $\Delta_1 c = c \otimes a + a^* \otimes c$ and the counit is $\epsilon_1(a) = 1$ and $\epsilon_1(c) = 0$.

Proposition 3.7 *There exists a coassociative coalgebra which is achiral entangled to the usual coassociative coalgebra part of $SU_q(2)$.*

Proof: Define the (left) coproduct $\tilde{\Delta}_1$ by the $*$ -linear map $a \mapsto \tilde{\Delta}_1 a := c^* \otimes a + a \otimes c$ and $c \mapsto \tilde{\Delta}_1 c := c \otimes c - q^{-1}a^* \otimes a$. This coproduct is coassociative. Moreover, we have $(\tilde{\Delta}_1 \otimes id)\Delta_1 = (id \otimes \Delta_1)\tilde{\Delta}_1$ and $(\Delta_1 \otimes id)\tilde{\Delta}_1 = (id \otimes \tilde{\Delta}_1)\Delta_1$. For instance, $(\Delta_1 \otimes id)\tilde{\Delta}_1 a = c^* \otimes a^* \otimes a + a \otimes c^* \otimes a + a \otimes a \otimes c - qc^* \otimes c \otimes c$ and $(id \otimes \tilde{\Delta}_1)\Delta_1 a = a \otimes c^* \otimes a + a \otimes a \otimes c - q(c^* \otimes c \otimes c - q^{-1}c^* \otimes a^* \otimes a)$, and so forth. The $*$ -linear map $\tilde{\epsilon}_1$ is defined by $c \mapsto 1$ and $a \mapsto 0$ is the (left) counit. $\square$

Remark: The usual algebraic relations of $SU_q(2)$ do not embed the new coproduct and the new counit into homomorphisms.

3.2 Splitting the coassociative coproducts and achiral L -coalgebras

The example of dendriform algebras ⁷ shows that there exists associative algebras whose associative product can split into two (a priori non associative) products. In the case of achiral L -coalgebras, we have also a splitting of a coassociative coproduct into two coassociative ones. Let us see how it works.

Recall that an achiral L -coalgebra $(L, \Delta, \tilde{\Delta})$ has two coproducts Δ and $\tilde{\Delta}$ which verify the axioms:

1. Δ and $\tilde{\Delta}$ are coassociative.
2. $(\tilde{\Delta} \otimes id)\Delta = (id \otimes \Delta)\tilde{\Delta}$.
3. $(\Delta \otimes id)\tilde{\Delta} = (id \otimes \tilde{\Delta})\Delta$,

i.e. the axioms are globally invariant by the permutation $\Delta \leftrightarrow \tilde{\Delta}$.

Proposition 3.8 *Suppose L is an achiral L -coalgebra. Let $u, v, w, z \in k$. These axioms are invariant under the transformation $\begin{pmatrix} \Delta \\ \tilde{\Delta} \end{pmatrix} \mapsto \begin{pmatrix} u & v \\ w & z \end{pmatrix} \begin{pmatrix} \Delta \\ \tilde{\Delta} \end{pmatrix}$. For instance, the coproduct $\Delta' := u\Delta + v\tilde{\Delta}$ is still coassociative.*

Proof: Straightforward. □

4 Tiling the $(n^2, 1)$ -De Bruijn co-dialgebra with n coassociative coalgebras

We can generalize the previous procedure to any coassociative coalgebras whose geometric supports are obtained by line-extension of the $(n, 1)$ -De Bruijn graphs viewed as geometric supports of Markov co-dialgebras. Fix $n > 1$, the number of vertices of the $(n, 1)$ -De Bruijn graph, denoted by U_i , $i = 1, \dots, n$. We have seen in proposition 2.10 that the line-extension of such a graph yields a geometric support of a coassociative coalgebra whose coproduct is denoted by Δ . The coproduct was recovered by computing $(U \bar{\otimes} U)$, where U_{ij} is the vertex associated with the arrow going from the vertex U_i to U_j in the $(n, 1)$ -De Bruijn graph. Denote by p , the shift which maps j into $j + 1 \pmod n$, for all $j = 1, \dots, n$. Let α be an integer equal to $0, \dots, n - 1$. We denote by $\mathcal{P}^\alpha(U)$, (resp. $\mathcal{P}^{-\alpha}(U)$), the matrix obtained from U by letting the shift p^α , (resp. $p^{-\alpha}$) acts on the columns of U , i.e. $\mathcal{P}^\alpha(U)_{ij} := U_{ip^\alpha(j)}$ and $\mathcal{P}^{-\alpha}(U)_{ij} := U_{ip^{-\alpha}(j)}$.

Lemma 4.1 *Let A, B be two n by n matrices and α be an integer equal to $0, \dots, n - 1$. We get $\mathcal{P}^\alpha(AB) = A\mathcal{P}^\alpha(B)$ and $\mathcal{P}^{-\alpha}(AB) = A\mathcal{P}^{-\alpha}(B)$. Therefore, we obtain, $\mathcal{P}^\alpha(U)U = \mathcal{P}^\alpha(\mathcal{P}^\alpha(U)\mathcal{P}^{-\alpha}(U))$.*

Proof: Notice that $\mathcal{P}^\alpha(AB)_{ij} = (AB)_{ip^\alpha(j)} = \sum_k A_{ik} B_{kp^\alpha(j)} = (A\mathcal{P}^\alpha(B))_{ij}$, which proved the first equality. The sequel is now straightforward. □

Remark: This lemma is also valid by replacing the usual product by $\bar{\otimes}$.

⁷See the introduction or [8].

Definition 4.2 [Coproducts] As $\mathcal{P}^0 = id$, we rename the usual coproduct Δ by $\Delta_{[0]}$. Its explicit definition, as we have seen, is closely related to the matrix U . We define also the coproducts $\Delta_{[\alpha]}$, for all α , by $\Delta_{[\alpha]}U_{ij} = (\mathcal{P}^\alpha(\mathcal{P}^\alpha(U) \bar{\otimes} \mathcal{P}^{-\alpha}(U)))_{ij} = (\mathcal{P}^\alpha(U)U)_{ij}$. Denote by $(\mathcal{F}_n, \Delta_{[\alpha]})$, $0 \leq \alpha \leq n-1$, the coalgebra so obtained. (Observe that $\mathcal{F}_2 := \mathcal{F}$.)

Theorem 4.3 *The new coproducts $\Delta_{[\alpha]}$ are coassociative. Moreover, for all $\alpha, \beta = 0, \dots, n-1$, $\Delta_{[\alpha]}$ and $\Delta_{[\beta]}$ obey the entanglement equation.*

Proof: We define for two matrices A, B the following product $A *_\alpha B = (\mathcal{P}^\alpha(\mathcal{P}^\alpha(A) \bar{\otimes} \mathcal{P}^{-\alpha}(B)))$. From the straightforward equality $A *_\alpha (B *_\alpha C) = (A *_\alpha B) *_\alpha C$, we obtain in the case where $A = B = C = U$, the coassociativity equation $(\Delta_{[\alpha]} \otimes id)\Delta_{[\alpha]}U_{ij} = (id \otimes \Delta_{[\alpha]})\Delta_{[\alpha]}U_{ij}$. To prove that two coproducts obey the entanglement equation, we have to show that $(\Delta_{[\alpha]} \otimes id)\Delta_{[\beta]}U_{ij} = (id \otimes \Delta_{[\beta]})\Delta_{[\alpha]}U_{ij}$, that is $U *_\alpha (U *_\beta U) = (U *_\alpha U) *_\beta U$, which is also straightforward. $\square$

Remark: We have showed that in the case of a coassociative coalgebra obtained by line-extension of the $(n, 1)$ -De Bruijn graphs, we can construct others coassociative coalgebras whose coproducts verify the entanglement equation. Precisely, the coassociative coalgebras $(\mathcal{F}_n, \Delta_{[\alpha]})$ and $(\mathcal{F}_n, \Delta_{[\beta]})$, $0 \leq \alpha, \beta \leq n-1$, are entangled, the entanglement being achiral. Therefore, the k -vector space $(\mathcal{F}_n, \Delta_{[0]}, \dots, \Delta_{[n-1]})$ is achiral⁸.

Proposition 4.4 *Let $(C, \Delta_{[0]}, \dots, \Delta_{[n-1]})$ be a k -vector space equipped with n linear maps $\Delta_{[i]} : C \rightarrow C^{\otimes 2}$, $i = 0, \dots, n-1$, such that:*

$$(\Delta_i \otimes id)\Delta_j = (id \otimes \Delta_j)\Delta_i, \quad i, j = 0, \dots, n-1.$$

For $i = 0, \dots, n-1$, set x_i the vector equal to ${}^t(0, \dots, 0, \Delta_i, 0, \dots, 0)$ and fix $Z \in M_n(k)$, a n by n matrix. Then $(C, \Delta'_{[0]} := Zx_0, \dots, \Delta'_{[n-1]} := Zx_{n-1})$ verify also:

$$(\Delta'_i \otimes id)\Delta'_j = (id \otimes \Delta'_j)\Delta'_i, \quad i, j = 0, \dots, n-1.$$

Proof: Straightforward. $\square$

Remark: Observe that this theorem allow us to produce splittings of associative laws into associative ones.

Remark: Applying this proposition to the tiling of the $(3, 1)$ -De Bruijn graph gives examples of cubical cotrialgebras, i.e. a coalgebra equipped with 3 coproducts Δ_i , $i = 0, 1, 2$, such that:

$$(\Delta_i \otimes id)\Delta_j = (id \otimes \Delta_j)\Delta_i, \quad i, j = 0, 1, 2.$$

The notion of cubical trialgebra is defined in [10]. We recover easily cubical trialgebras from cubical cotrialgebras by considering the convolution products. The operad *Tricub* on one generator associated with cubical trialgebra is Koszul and self-dual. The operad associated with the so-called hypercube n -algebra, i.e. a k -vector space equipped with n products verifying:

$$(x \bullet_i y) \bullet_j z = x \bullet_i (y \bullet_j z), \quad x, y, z \in A, \quad i, j = 0, \dots, n-1,$$

⁸In [5], such algebraic objects will be called coassociative manifolds.

is conjectured to be Kozul and self-dual (observe that there are n^2 operations and two possible choices of parentheses, thus $2n^2 - n^2 = n^2$).

Remark: The counit $\epsilon_{[\alpha]}$, associated with the coassociative coproduct $\Delta_{[\alpha]}$, is obviously defined by $U_{ip^\alpha(i)} \mapsto 1$ and $U_{ip^\alpha(j)} \mapsto 0$ if $i \neq j$.

Theorem 4.5 *The intersection of the arrow sets of the geometric supports of $(\mathcal{F}_n, \Delta_{[\alpha]})$, $0 \leq \alpha \leq n-1$, are empty, i.e. $Gr((\mathcal{F}_n, \Delta_{[\alpha]}))_1 \cap Gr((\mathcal{F}_n, \Delta_{[\beta]}))_1 = \emptyset$, $0 \leq \alpha, \beta \leq n-1$ and $\alpha \neq \beta$. Gluing them yields the $(n^2, 1)$ -De Bruijn directed graph, i.e. $\bigcup_{0 \leq \alpha \leq n-1} Gr((\mathcal{F}_n, \Delta_{[\alpha]})) = D_{(n^2, 1)}$.*

Proof: We call $(U_{ij})_{i,j=1,\dots,n}$, the vertices of the $(n^2, 1)$ -De Bruijn graph. Let us prove that the gluing of all the geometric supports of $(\mathcal{F}_n, \Delta_{[\alpha]})$, $0 \leq \alpha \leq n-1$, yields the $(n^2, 1)$ -De Bruijn directed graph. Every arrow of the $(n^2, 1)$ -De Bruijn directed graph can be described by $U_{ik} \otimes U_{lj}$, with $k, l, i, j = 1, \dots, n$. As the shift is one-to-one, there exists a unique integer α such that $p^\alpha(l) = k$ i.e. $U_{ik} \otimes U_{lj} = U_{ip^\alpha(l)} \otimes U_{lj}$, i.e. this arrow belongs to the definition of the coproduct $\Delta_{[\alpha]}$, see lemma 4.1. Therefore, the $(n^2, 1)$ -De Bruijn graph is included into the gluing of the geometric support of the n coassociative coalgebras. The reversal is obvious.

Fix α and β two different integers. As the shift $p^{\beta-\alpha}$ has no fixed point, no arrow defined in the coproduct $\Delta_{[\alpha]}$ is present in the definition of the coproduct of $\Delta_{[\beta]}$. Therefore, the intersection of arrow sets of $Gr(\mathcal{F}_n, \Delta_{[\alpha]})$ and $Gr(\mathcal{F}_n, \Delta_{[\beta]})$ is empty. $\square$

Remark: Via their geometric supports, we get:

$(n, 1)$ -De Bruijn co-dialgebra $\xrightarrow{\text{Line-extension}} n$ coassociative coalgebras $\xrightarrow{9 \text{ Gluing}} (n^2, 1)$ -De Bruijn co-dialgebra.

We have yielded a tiling of the $(n^2, 1)$ -De Bruijn graph into n coassociative coalgebras, each $(\mathcal{F}_n, \Delta_{[\alpha]}, \Delta_{[\beta]})$, $0 \leq \alpha, \beta \leq n-1$, being an achiral coassociative L -coalgebra. The case $n = 1$ is trivial. Indeed, the $(1, 1)$ -De Bruijn graph and its line-extension are loops and a loop is coassociative, since it is of the form $x \mapsto x \otimes x$.

Remark: Fix $n \geq 1$. The link between the Markovian coproducts of the coassociative codialgebra $(k(D_{(n,1)})_0, \Delta_M, \tilde{\Delta}_M)$, associated with the $(n, 1)$ -De Bruijn graph and the coassociative coproduct of $(\mathcal{F}_n, \Delta_{[0]})$ can be seen as follows. Recall that $D_{(n,1)}$ is the $(n, 1)$ -De Bruijn graph, with vertex set $(D_{(n,1)})_0 := \{U_i; 1 \leq i \leq n\}$. Consider its free k -vector space $k(D_{(n,1)})_0$. Its Markovian coproducts verify $\Delta_M U_i := U_i \otimes \sum_k U_k$ and $\tilde{\Delta}_M U_i := \sum_k U_k \otimes U_i$. Set for all $i, j = 0, \dots, n-1$, $U_{ij} := U_i \otimes U_j$. Define the map $\boxtimes : k(D_{(n,1)})_0^{\otimes 2} \times k(D_{(n,1)})_0^{\otimes 2} \rightarrow k(D_{(n,1)})_0^{\otimes 2} \otimes k(D_{(n,1)})_0^{\otimes 2}$ such that $(z_1 \otimes z_2) \boxtimes (z_3 \otimes z_4) := (id \otimes \Psi \otimes id)(z_1 \otimes z_2 \otimes z_3 \otimes z_4)$, where $\Psi(z_i \otimes z_j) = z_i \otimes z_i$ if $i = j$, and 0 otherwise. Observe that:

$$\Delta_M(U_i) \boxtimes \tilde{\Delta}_M(U_j) := \Delta_{[0]} U_{ij}, \quad i, j = 0, \dots, n-1.$$

As in L -coalgebra theory, the notion of algebraic product is not assumed, we end this section by a proposition on the Leibniz coderivation and an example applied on $\mathcal{F}$.

⁹entangled by the achiral entanglement equation.

Definition 4.6 [Coderivation] Let C be a coassociative coalgebra with coproduct Δ . A linear map $D : C \rightarrow C$ is called a (Leibniz) *coderivation* with respect to the coproduct Δ iff it verifies:

$$\Delta D = (id \otimes D)\Delta + (D \otimes id)\Delta.$$

Proposition 4.7 Let C be a coassociative coalgebra with coproduct Δ , such that for all the elements $U_{ij} \in C$, $\Delta U_{ij} = \sum_{k=1}^n U_{ik} \otimes U_{kj}$, where n is a fixed integer. Define the linear map D by $U_{ij} \mapsto \sum_k^n U_{kj} - U_{ik}$. Then D is a Leibniz coderivation.

Proof: Fix U_{ij} . We have $(id \otimes D)\Delta + (D \otimes id)\Delta(U_{ij}) = \sum_{k,l=1}^n U_{lk} \otimes U_{kj} - U_{ik} \otimes U_{kl}$ which is equal to $\Delta(\sum_{l=1}^n U_{lj} - U_{il})$. $\square$

Example 4.8 [A coderivation for $(\mathcal{F}, \Delta, \tilde{\Delta})$]

For $(\mathcal{F}, \Delta)$, we get $D(a) = c - b = -D(d)$, $D(b) = d - a = -D(c)$. As $(\mathcal{F}, \Delta, \tilde{\Delta})$ is a L -coalgebra, we yield also a coderivation $\tilde{D}$ for $(\mathcal{F}, \tilde{\Delta})$. Define $\tilde{D} = D$. A straightforward computation shows that D is also a coderivation with respect to the coproduct $\tilde{\Delta}$.

5 Conclusion

The first main result obtained in this paper is the possibility to construct from directed graphs, families of L -cocommutative coassociative co-dialgebras and therefore, via convolution products families of associative dialgebras. The second main result is the possibility to recover from the line-extension of the geometric supports of these coassociative co-dialgebras, known coassociative coalgebras and in the case of the $(n^2, 1)$ -De Bruijn graphs, $n > 0$, to obtain a tiling of these Markovian objects by n (geometric supports of) coassociative coalgebras. An important notion, called the achirality has been put forward. We have shown that actions of $M_n(k)$ on the coproducts defining the tiling of the $(n^2, 1)$ -De Bruijn graph let globally invariant the relations between them. We gave consequences of such tilings and found examples of cubical trialgebras and more generally, examples of hypercube n -algebras. In addition, this allowed us to construct associative laws which split into several associative ones.

This paper has been pursued in [5]. In [5], the notion of codipterous coalgebras and pre-dendriform coalgebras¹⁰ are established. These spaces constructed from coassociative coalgebra theory extend the notions developed so far and are the elementary boxes of coassociative manifolds. Via these notions, we construct Poisson algebras, dendriform algebras, associative dialgebras (which are not Markovian), associative trialgebras [10]. Notably, the tilings constructed so far will yield examples of coassociative manifolds [5].

Acknowledgments: The author wishes to thank Dimitri Petritis for useful discussions and fruitful advice for the redaction of this paper and to S. Severini for pointing him the precise definition of the De Bruijn graphs.

¹⁰These notions have been discovered by J-L Loday and M. Ronco [9] and rediscovered independently, via graph theory by the author [1].

References

- [1] Private communication.
- [2] A. CONNES and D. KREIMER. Hopf algebras, renormalization and non commutative geometry. *eprint arXiv:hep-th/9912092*, *Commun. Math. Phys.*, 199:203–242, 1998.
- [3] D. KREIMER. Chen’s iterated integral represents the operator product expansion. *eprint arXiv:hep-th/9901099*, *Adv. Theor. Math. Phys.*, 3.3, 1999.
- [4] Ph. LEROUX. Coassociativity breaking and oriented graphs. *eprint arXiv:math.QA/0204342*.
- [5] Ph. LEROUX. From entangled codipterous coalgebras to coassociative manifolds. *eprint arXiv:math.QA/0301080*.
- [6] Ph. LEROUX. On representations of braid groups determined by directed graphs. *eprint arXiv:math.QA/0210260*.
- [7] Ph. LEROUX. Periodic orbits, coassociative grammar and quantum random walk on $\mathbb{Z}$. *eprint arXiv:quant-ph/0209100*.
- [8] J-L. LODAY. Dialgebras. *eprint arXiv:math.QA/0102053*, in *Dialgebras and related operads, Lecture Notes in Mathematics*, 1763:7–66, 2001.
- [9] J-L. LODAY and M. RONCO. In preparation.
- [10] J-L. LODAY and M. RONCO. Trialgebras and families of polytopes. *eprint arXiv:math.QA/0205043*.
- [11] K.H. ROSEN, J.G. MICHAELS, J.L. GROSS, J.W. GROSSMAN, and D.R. SHIER. *Handbook of Discrete and Combinatorial Mathematics*. 2000.